GASTROENTEROLOGY OF SAN MARCOS/NEW BRAUNFELS

Texas Digestive Disease Consultants

Diplomates of the American Board of Internal Medicine and Subspecialty Board of Gastroenterology

W. Kern Deschner, M.D., F.A.C.G. Edwards F. Coles, M.D., F.A.C.P., F.A.C.G. Thomas Kepczyk, M.D., F.A.C.P., F.A.C.G. Carlos A Neghme, M.D., F.A.C.G. Shannon Marek, M.D. Steven Pearce, M.D. John Cluley, M.D. Lori Hale, PA-C, MPAS

COLONOSCOPY Trilyte

PLEASE READ ALL INSTRUCTIONS CAREFULLY!

Vou have been schoduled for a colonoscopy on

Tou have been scheduled for a	cololloscopy off	aı	•
Please arrive at facility by	•		
	- · · · · · · · · · · · · · · · · · · ·	****	******

- o Bring your Insurance Card and payment for necessary co-pays
- o Wear loose-fitting comfortable clothing. NO lipstick, contact lenses, makeup.
- Leave Jewelry at home
- o Bring phone and fax numbers of referring physicians you would like reports sent to
- YOU MUST HAVE A RESPONSIBLE ADULT TO DRIVE YOU HOME
 - Your driver must stay during your procedure. Plan for someone to be with you 4-6 hours after your procedure as well.
 - o Your procedure WILL BE CANCELLED if you do not have a driver to take you home.

MEDICATION GUIDELINES



- You may take your other necessary medications as you usually would
- On day of the exam, NO diuretics should be taken (Lasix, Aldactone, HCTZ, etc)
- Diabetic medications require special instructions
- BRING A LIST OF YOUR CURRENT MEDICATIONS (including over the counter meds)
- If you are a female, you must notify someone if there is any possibility you maybe pregnant.

5 days prior to your procedure (

- STOP TAKING: Iron pills, Coumadin, Plavix, Aggrenox, Pletal, Effient, Pradexa, Xarelto, Eliquis etc. (unless otherwise instructed)
- Avoid eating any <u>raw</u> fruits, vegetables or nuts(COOKED OR CANNED VEGETABLES AND FRUITS ARE ALLOWED)

GASTROENTEROLOGY OF SAN MARCOS/NEW BRAUNFELS

Texas Digestive Disease Consultants

Diplomates of the American Board of Internal Medicine and Subspecialty Board of Gastroenterology

W. Kern Deschner, M.D., F.A.C.G.
Edwards F. Coles, M.D., F.A.C.P., F.A.C.G.
Thomas Kepczyk, M.D., F.A.C.P., F.A.C.G.
Carlos A Neghme, M.D., F.A.C.G.

Shannon Marek, M.D.
Steven Pearce, M.D.
John Cluley, M.D.
Lori Hale, PA-C, MPAS

IF YOU HAVE NOT HAD A BOWEL MOVEMENT 1-2 DAYS PRIOR TO YOUR EXAM, TAKE 1 BOTTLE OF MAGNESIUM CITRATE (which is sold over the counter) AT 8PM ON .

day prior to your procedure ()

- FOR BREAKFAST: You may have a full liquid diet (milk, plain oatmeal, plain yogurt, pudding, nut free ice cream, cream soups)
- REST OF THE DAY STARTING AT 12 NOON: CLEAR LIQUID DIET ONLY. NO SOLID FOODS
 - Strained Fruit Juices without pulp (apple, White Grape, Orange or Lemonade) No red/purple juices.
 - Water (drink at least 64 ounces through the course of the day)
 - Clear broth or bouillon, coffee or tea, (without milk or non-dairy creamer)
 - All of the following as long as they are NOT colored RED or PURPLE
 - Gatorade- Snapple
 - Carbonated and non-carbonated soft drinks (7-up, sprite, ginger ale)
 - Kool-Aid (or other fruit flavored drinks)
 - Plain Jell-O (preferably clear or yellow, without added fruits or toppings)
- AT 6:30PM: Take the first dose of Reglan (generic name: Metoclopramide) to prevent nausea
- AT 7:00PM: Take first dose of TRILYTE laxative
 - o You will have to mix the entire gallon as instructed on prescription.
 - Add lukewarm drinking water to the top line of the container. Mix to dissolve. If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.
 - o (OPTIONAL) You can mix Crystal Light (no red flavors) with the solution for flavor
 - o You will need to drink ½ of the gallon. You will need to drink one 8 ounce glass every 15 minutes until ½ of the gallon is consumed. Place the remaining ½ gallon of solution in the refrigerator for your 2nd dose.
 - o (OPTIONAL) Flushable moist wipes may prevent anal irritation from excessive wiping with the prep

Day of your procedure

- 6.5 HOURS PRIOR (at
-): Take the 2nd dose of Reglan (generic name: Metoclopramide) to prevent nausea
- 6 HOURS PRIOR (at
-): Take second dose of TRILYTE laxative
- o You will drink the remaining ½ gallon of solution. Following instructions as above.
- 4 HOURS PRIOR (at): NO FURTHER LIQUIDS BY MOUTH(no gum,candy,ice or water) (except a sip of water with blood pressure medication only) Hold all other medications until after your procedure.

GASTROENTEROLOGY OF SAN MARCOS/NEW BRAUNFELS

Texas Digestive Disease Consultants

Diplomates of the American Board of Internal Medicine and Subspecialty Board of Gastroenterology

W. Kern Deschner, M.D., F.A.C.G.
Edwards F. Coles, M.D., F.A.C.P., F.A.C.G.
Thomas Kepczyk, M.D., F.A.C.P., F.A.C.G.
Carlos A Neghme, M.D., F.A.C.G.

Shannon Marek, M.D. Steven Pearce, M.D. John Cluley, M.D. Lori Hale, PA-C, MPAS

******* If you eat or drink within 4 hours of the procedure, we may have to cancel the procedure.********
If you have any questions please call: Barbara ext 2205 or Julie ext 2228
YOU ARE SCHEDULED AT THE FOLLOWING:
Central Texas Medical Center (call 512-753-3665) 1-2 days prior to exam 1301 Wonder World Drive, San Marcos, TX. 78666
Christus Santa Rosa Out Patient Surgery-San Marcos Enter through "Main Entrance" of Central Texas Medical Center (call 830-643-8600) 1-2 days prior to exam 1301 Wonder World Drive Suite 105, San Marcos, TX. 78666
Christus Santa Rosa Out Patient Surgery-New Braunfels (call 830-643-8600)1-2 days prior to exam 1738 E. Common Street, New Braunfels, TX 78130 (YOU ARE SCHEDULED AT SURGERY CENTER NOT HOSPITAL)
Seton Edgar B. Davis Admission Office (call 830-875-7000)1-2 days prior to exam 130 Hays Street, Luling, TX. 78648
Resolute Hospital (830) 500-6000 (No pre-registration required) 555 Creekside Way, New Braunfels, TX 78130
 For questions regarding anesthesia call 1-866-647-3631 For questions regarding physicians bill call 214-623-6230